

大同大學 97 學年度研究所碩士班入學考試試題

考試科目：統計學

所別：資訊經營研究所

第 1/3 頁

註：本次考試 不可以參考自己的書籍及筆記； 不可以使用字典； 可以使用計算器。

1. Selection (3 points for each question)

- (1) In a questionnaire, respondents are asked to mark their gender as male or female. Gender is an example of the
 - a. ordinal scale
 - b. nominal scale
 - c. ratio scale
 - d. interval scale
- (2) Some hotels ask their guests to rate the hotel's services as excellent, very good, good, and poor. This is an example of the
 - a. ordinal scale
 - b. ratio scale
 - c. nominal scale
 - d. interval scale
- (3) Temperature is an example of a variable that uses
 - a. the ratio scale
 - b. the interval scale
 - c. the ordinal scale
 - d. either the ratio or the ordinal scale
- (4) Quantitative data refers to data obtained with a(n)
 - a. ordinal scale
 - b. nominal scale
 - c. either interval or ratio scale
 - d. only interval scale
- (5) A frequency distribution is a tabular summary of data showing the
 - a. fraction of items in several classes
 - b. percentage of items in several classes
 - c. relative percentage of items in several classes
 - d. number of items in several classes
- (6) A situation in which conclusions based upon aggregated crosstabulation are different from unaggregated crosstabulation is known as
 - a. wrong crosstabulation
 - b. Simpson's rule
 - c. Simpson's paradox
 - d. aggregated crosstabulation
- (7) Data that provide labels or names for categories of like items are known as
 - a. qualitative data
 - b. quantitative data
 - c. label data
 - d. category data
- (8) The most common graphical presentation of quantitative data is a
 - a. histogram
 - b. bar graph
 - c. relative frequency
 - d. pie chart
- (9) When computing the mean of a set of values , the value of
 - a. can never be zero
 - b. can never be negative
 - c. must always be positive
 - d. can be any value

<背面繼續>

大同大學 97 學年度研究所碩士班入學考試試題

考試科目：統計學

所別：資訊經營研究所

第 2/3 頁

註：本次考試 不可以參考自己的書籍及筆記； 不可以使用字典； 可以使用計算器。

<接前頁>

- (10) The interquartile range is
- the 50th percentile
 - another name for the variance
 - the difference between the largest and smallest values
 - the difference between the third quartile and the first quartile
2. The time needed to complete a final examination in a particular college course is normally distributed with a mean of 80 minutes and a standard deviation of 10 minutes. Answer the following questions. (5 points for each question)
- What is the probability of completing the exam in one hour or less?
 - What is the probability that a student will complete the exam in more than 60 minutes but less than 75 minutes?
 - Assume that the class has 60 students and that the examination period is 90 minutes in length. How many students do you expect will be unable to complete the exam in the allotted time?
3. According to Barron's 1998 Primary Reader Survey, the average annual number of investment transactions for a subscriber is 30. Suppose the number of transactions in a year follows the Poisson probability distribution. (5 points for each question)
- Show the probability distribution for the time between investment transactions.
 - What is the probability of no transactions during the month of January for a particular subscriber?
 - What is the probability that the next transaction will occur within the next half month for a particular subscriber?
4. A Taipei City government survey found that households tend to spend an average of NT\$17,576 for food and beverage during Chinese Lunar New Year. Assume that the survey included 600 households and the standard deviation was NT\$4,740. (5 points for each question)
- With 95% confidence interval, what is the margin of error?
 - What is the 95% confidence interval estimate of the population mean?
5. The mean number of hours during a work week for the population of workers was reported to be 44.75 hours. Suppose we take a current sample of workers to see whether the mean number of hours over a work week has changed from the previously reported 44.75 hours. (5 points for each question)
- State the hypotheses that will help us determine whether a change occurred in the mean length of work week?
 - Suppose a current sample of 111 workers provided a sample mean of 43.95 hours. Using a population standard deviation $\sigma = 5.48$ hours, what is the p -value?
 - At $\alpha = .05$, can the null hypothesis be rejected? What is your conclusion?
 - Repeat the preceding hypothesis test using the critical value approach.
6. *Young Adult* magazine states the following hypotheses about the mean age of its subscribers. (5 points for each question)
- $$H_0 : \mu = 28$$
- $$H_a : \mu \neq 28$$
- What would it mean to make a Type II error in this situation?
 - If the manager conducting the test will permit a .15 probability of making a Type II error when the true mean age is 29, what sample size should be selected? Assume $\sigma = 6$ and a .05 level of significance.

大同大學 97 學年度研究所碩士班入學考試試題


考試科目：統計學

所別：資訊經營研究所

第 3/3 頁

註：本次考試 不可以參考自己的書籍及筆記； 不可以使用字典； 可以使用計算器。

TABLE 1. STANDARD NORMAL DISTRIBUTION




Entries in the table give the area under the curve between the mean and z standard deviations above the mean. For example, for $z = 1.25$ the area under the curve between the mean and z is .3944.

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0753
.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2517	.2549
.7	.2580	.2611	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	.4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.4987	.4987	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990

TABLE 6. VALUES OF e^{-z}

z	e^{-z}	z	e^{-z}	z	e^{-z}
.00	1.0000	2.00	.1353	4.00	.0183
.05	.9512	2.05	.1287	4.05	.0174
.10	.9048	2.10	.1225	4.10	.0166
.15	.8607	2.15	.1165	4.15	.0158
.20	.8187	2.20	.1108	4.20	.0150
.25	.7788	2.25	.1054	4.25	.0143
.30	.7408	2.30	.1003	4.30	.0136
.35	.7047	2.35	.0954	4.35	.0129
.40	.6703	2.40	.0907	4.40	.0123
.45	.6376	2.45	.0863	4.45	.0117
.50	.6065	2.50	.0821	4.50	.0111
.55	.5769	2.55	.0781	4.55	.0106
.60	.5488	2.60	.0743	4.60	.0101
.65	.5220	2.65	.0707	4.65	.0096
.70	.4966	2.70	.0672	4.70	.0091
.75	.4724	2.75	.0639	4.75	.0087
.80	.4493	2.80	.0608	4.80	.0082
.85	.4274	2.85	.0578	4.85	.0078
.90	.4066	2.90	.0550	4.90	.0074
.95	.3867	2.95	.0523	4.95	.0071
1.00	.3679	3.00	.0498	5.00	.0067
1.05	.3499	3.05	.0474	6.00	.0025
1.10	.3329	3.10	.0450	7.00	.0009
1.15	.3166	3.15	.0429	8.00	.000335
1.20	.3012	3.20	.0408	9.00	.000123
				10.00	.000045
1.25	.2865	3.25	.0388		
1.30	.2725	3.30	.0369		
1.35	.2592	3.35	.0351		
1.40	.2466	3.40	.0334		
1.45	.2346	3.45	.0317		
1.50	.2231	3.50	.0302		
1.55	.2122	3.55	.0287		
1.60	.2019	3.60	.0273		
1.65	.1920	3.65	.0260		
1.70	.1827	3.70	.0247		
1.75	.1738	3.75	.0235		
1.80	.1653	3.80	.0224		
1.85	.1572	3.85	.0213		
1.90	.1496	3.90	.0202		
1.95	.1423	3.95	.0193		

TABLE 2. t DISTRIBUTION



Entries in the table give t values for an area or probability in the upper tail of the t distribution. For example, with 10 degrees of freedom and a .05 area in the upper tail, $t_{.05} = 1.812$.

Degrees of Freedom	Area in Upper Tail					
	.20	.10	.05	.025	.01	.005
80	.846	1.292	1.664	1.990	2.374	2.639
81	.846	1.292	1.664	1.990	2.373	2.638
82	.846	1.292	1.664	1.989	2.373	2.637
83	.846	1.292	1.663	1.989	2.372	2.636
84	.846	1.292	1.663	1.989	2.372	2.636
85	.846	1.292	1.663	1.988	2.371	2.635
86	.846	1.291	1.663	1.988	2.370	2.634
87	.846	1.291	1.663	1.988	2.370	2.634
88	.846	1.291	1.662	1.987	2.369	2.633
89	.846	1.291	1.662	1.987	2.369	2.632
90	.846	1.291	1.662	1.987	2.368	2.632
91	.846	1.291	1.662	1.986	2.368	2.631
92	.846	1.291	1.662	1.986	2.368	2.630
93	.846	1.291	1.661	1.986	2.367	2.630
94	.845	1.291	1.661	1.986	2.367	2.629
95	.845	1.291	1.661	1.985	2.366	2.629
96	.845	1.290	1.661	1.985	2.366	2.628
97	.845	1.290	1.661	1.985	2.365	2.627
98	.845	1.290	1.661	1.984	2.365	2.627
99	.845	1.290	1.660	1.984	2.364	2.626
100	.845	1.290	1.660	1.984	2.364	2.626
∞	.842	1.282	1.645	1.960	2.326	2.576